


SAMPLE PRESCRIBER'S ORDERS

This is an example of a pre-printed order that may be used in the provision of iOAT.



Date and time	TITRATION ORDERS FOR HIGH-DOSE HYDROMORPHONE (Items with checkboxes must be selected to be ordered)
	<p style="text-align: center;">These orders are applicable to clients who are starting on HYDROmorphone treatment</p> <p>INDICATION: Opioid use disorder</p> <p>MEDICATION: HYDROmorphone 50 mg/mL injection for the following orders for ALL routes of administration (IM/IV)</p> <p>Qualified health professional to assess for signs and symptoms of opioid sedation and withdrawal before and after each dose using Pasero Opioid-induced Sedation Scale (POSS) If POSS 3 or 4, notify prescriber immediately and hold next dose</p> <ul style="list-style-type: none"> <input type="checkbox"/> Qualified health professional to assess for signs and symptoms of stimulant use before each dose and notify prescriber if the following symptoms are noted: _____ <input type="checkbox"/> Qualified health professional to assess for alcohol use with breathalyzer before each dose. Hold dose and notify prescriber if breathalyzer reading is: _____ <ul style="list-style-type: none"> • Rapid urine drug testing PRN • Breathalyzer PRN • Qualified health professional to notify prescriber if client misses 2 or more sessions or declines second dose • Qualified health professional to notify prescriber about titration by 3pm on Day 2 in order for prescriber to write maintenance prescription • Round all doses UP to nearest 5mg <p>Day 1, Session 1:</p> <p>Dose 1: HYDROmorphone 10mg (ten milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose</p> <p>Dose 2: HYDROmorphone 10mg (ten milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose Qualified health professional to advise client of timing for return for Session 2</p> <p style="text-align: center;">Do not exceed 20mg (twenty milligrams) of HYDROmorphone IM/IV for Day 1, Session 1</p> <p>Day 1, Session 2:</p> <p>Dose 1: HYDROmorphone 20mg (twenty milligrams) IV/IM for client self-administration x 1 dose Monitor for signs and symptoms of intoxication for 15 minutes post-dose</p> <p>Dose 2: HYDROmorphone 10mg (ten milligrams) IV/IM for client self-administration x 1 dose</p>

THIS IS A SAMPLE PRE-PRINTED ORDER ADAPTED FROM CROSSTOWN CLINIC IN VANCOUVER, BC.
 THIS FORM IS NOT MEANT FOR CLINICAL USE.

Monitor for signs and symptoms of intoxication for 15 minutes post-dose
Qualified health professional to advise client of timing for return for Session 3

Do not exceed 30mg (thirty milligrams) of HYDROmorphine IM/IV for Day 1, Session 2

Day 1, Session 3:

Dose 1: HYDROmorphine 30mg (thirty milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Dose 2: HYDROmorphine 10mg (ten milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose
Qualified health professional to advise client of timing for return for Day 2

Do not exceed 40mg (forty milligrams) of HYDROmorphine IM/IV for Day 1, Session 3

Day 2, Session 1:

Dose 1: HYDROmorphine 40 mg (forty milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Dose 2: HYDROmorphine 10mg (ten milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose
Qualified health professional to advise client of timing for return for Session 2

Do not exceed 50mg (fifty milligrams) of HYDROmorphine IM/IV for Day 2, Session 1

Day 2, Session 2:

If earlier doses were well tolerated:

Dose 1: HYDROmorphine 50mg (fifty milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Dose 2: HYDROmorphine 10mg (ten milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose
Qualified health professional to advise client of timing for return for Session 3

Do not exceed 60mg (sixty milligrams) of HYDROmorphine IM/IV for Day 2, Session 2

Day 2, Session 3:

Dose 1: If earlier doses were well tolerated:
HYDROmorphine 60mg (sixty milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Dose 2: HYDROmorphine 10mg (ten milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose
Qualified health professional to advise client of timing for return for Day 3

Do not exceed 70mg (seventy milligrams) of HYDROmorphine IM/IV for Day 2, Session 3

Day 3, Session 1:

Dose 1: HYDROmorphine 70mg (seventy milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose

THIS IS A SAMPLE PRE-PRINTED ORDER ADAPTED FROM CROSSTOWN CLINIC IN VANCOUVER, BC.
THIS FORM IS NOT MEANT FOR CLINICAL USE.

Dose 2: HYDROmorphone 10mg (ten milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose
Qualified health professional to advise client of timing for return for Session 3

Do not exceed 80mg (eighty milligrams) of HYDROmorphone IM/IV for Day 3, Session 1

Day 3, Session 2:

Dose 1: HYDROmorphone 80mg (eighty milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose

Dose 2: HYDROmorphone 10mg (ten milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose
Qualified health professional to advise client of timing for return for Day 3

Do not exceed 90mg (ninety milligrams) of HYDROmorphone IM/IV for Day 3, Session 2

Day 3, Session 3:

Dose 1: HYDROmorphone 90mg (ninety milligrams) IV/IM for client self-administration x 1 dose
Monitor for signs and symptoms of intoxication for 15 minutes post-dose
Qualified health professional to advise client of timing for return to clinic now that titration is complete

Do not exceed 90mg (ninety milligrams) of HYDROmorphone IM/IV for Day 3, Session 3

THIS IS A SAMPLE PRE-PRINTED ORDER ADAPTED FROM CROSTOWN CLINIC IN VANCOUVER, BC.
THIS FORM IS NOT MEANT FOR CLINICAL USE.